

First Report of German Cockroaches (*Blattella germanica*) as Reservoirs of CTX-M-15 Extended-Spectrum- β -Lactamase- and OXA-48 Carbapenemase-Producing *Enterobacteriaceae* in Batna University Hospital, Algeria

Lotfi Loucif,^{a,b} Djamil Gacemi-Kirane,^c Zineb Cherak,^a Naima Chamal,^a Nadia Grainat,^d Jean-Marc Rolain^b

Laboratoire de Biotechnologie des Molécules Bioactives et de la Physiopathologie Cellulaire (LBMBPC), Faculté des Sciences de la Nature et de la Vie, Université de Batna 2, Batna, Algeria^a; Unité de Recherche sur les Maladies Infectieuses et Tropicales Émergentes (URMITE), UM 63, CNRS 7278, IRD 198, INSERM 1095, IHU Méditerranée Infection, Faculté de Médecine et de Pharmacie, Aix-Marseille-Université, Marseille, France^b; Département de Biochimie, Faculté des Sciences, Université Badji Mokhtar Annaba, Annaba, Algeria^c; Faculté de Médecine, Université de Batna 2, Batna, Algeria^d

Here we report the isolation of extended-spectrum β -lactamase (ESBL)- and carbapenemase-producing *Enterobacteriaceae* from German cockroaches caught in the burn unit of Batna University Hospital in Algeria. Nine of 12 isolates harbored the *bla*_{CTX-M-15} ESBL gene. One *Enterobacter cloacae* isolate belonging to sequence type 528 coexpressed the *bla*_{OXA-48}, *bla*_{CTX-M-15}, and *bla*_{TEM} genes. Our findings indicate that cockroaches may be one of the most dangerous reservoirs for ESBL and carbapenemase producers in hospitals.

The emergence of multidrug-resistant *Enterobacteriaceae* (MDRE) and their dissemination in hospitals have become a major health concern worldwide (1). The production of β -lactamases such as extended-spectrum β -lactamase (ESBL) and carbapenemase remains the main mechanism of resistance in *Enterobacteriaceae* (2). Algeria is now features among the Mediterranean countries known to be affected by the spread of CTX-M-15 type ESBL and the emergence of the phantom menace OXA-48 type carbapenemase (3, 4). Indeed, the transmission and dissemination of MDRE in hospitals are usually related to hand contact (5) and poorly sterilized instruments (6). However, cockroaches which colonize these environments may also act as potential mechanical vectors of antibiotic-resistant bacteria (7). Cockroaches are among the most prevalent pests in hospitals, where they are attracted by moisture, food, and suitable temperatures (8, 9). The medical significance of cockroaches has been largely overlooked, but it has been proven that they can harbor a number of pathogenic microorganisms with different levels of antibiotic resistance (10). *Blattella germanica* is the most abundant cockroach species (11), and the external surfaces and excrement of these cockroaches can contaminate food, human habitats, and hospital equipment; therefore, these insects can present a threat to human health (12).

The aim of our study was to screen for broad-spectrum-cephalosporin- and carbapenem-resistant *Enterobacteriaceae* in *B. germanica* cockroaches collected from the burn unit of Batna University Hospital in Algeria and then to investigate the molecular support of ESBL and carbapenemase production.

In March 2015, 10 cockroaches were randomly captured from different parts of the burn unit, including the kitchen and the guard, treatment, and patient rooms, directly in sterile containers. Samples were immobilized by freezing at 0°C for 10 min (8). Each cockroach was soaked in 5 ml of Tween 80 solution at 0.05% and vortexed vigorously for 2 min. The resulting wash was used as an external-body-homogenate sample. To remove the external body contamination, cockroaches were immersed in bleach for 2 min, in sterile physiological saline for 2 min, and then in 70% ethanol

for 5 min (12). Then, each sample was washed with sterile physiological saline. Subsequently, the insect was soaked in a sterile bottle containing 5 ml of Tween 80 solution at 0.05% and then crushed inside using a sterile pestle. The triturate was then vortexed vigorously for 2 min. The resulting suspension was used as an internal-body-homogenate sample. Each sample (5 ml) was preenriched for 4 h in 5 ml of a double concentration of brain heart infusion (BHI) (BD: Becton, Dickinson and Company, France). Subsequently, the screening for β -lactam-resistant *Enterobacteriaceae* began with a selective enrichment step using two selective media: BHI supplemented with 64 mg/liter of vancomycin and 1 mg/liter of ertapenem and BHI supplemented with 64 mg/liter of vancomycin and 2 mg/liter of cefotaxime. After overnight incubation at 37°C, cultures were inoculated by streaking 100 μ l of the growth onto MacConkey agar plates (BD) with 64 mg/liter of vancomycin supplemented with 1 mg/liter of ertapenem or 2 mg/liter of cefotaxime, respectively. The obtained *Enterobacteriaceae* isolates were identified using an API 20E system (bioMérieux, France) and confirmed by matrix-assisted laser desorption ionization-time of flight mass spectrometry (13). Susceptibility testing was performed using a standard disk diffusion technique according to the recommendations of the Antibiotic Committee of the French Society for Microbiology (http://www.sfm-microbiologie.org/UserFiles/files/casfm/CASFM_EUCAST_V1_2015.pdf). ESBL detection was per-

Received 21 April 2016 Returned for modification 22 May 2016

Accepted 17 July 2016

Accepted manuscript posted online 25 July 2016

Citation Loucif L, Gacemi-Kirane D, Cherak Z, Chamal N, Grainat N, Rolain J-M. 2016. First report of German cockroaches (*Blattella germanica*) as reservoirs of CTX-M-15 extended-spectrum- β -lactamase- and OXA-48 carbapenemase-producing *Enterobacteriaceae* in Batna University Hospital, Algeria. *Antimicrob Agents Chemother* 60:6377–6380. doi:10.1128/AAC.00871-16.

Address correspondence to Jean-Marc Rolain, jean-marc.rolain@univ-amu.fr.

Copyright © 2016, American Society for Microbiology. All Rights Reserved.

formed using the double-disk synergy test (DDST) as previously described (14). The modified Hodge test (MHT) and the modified Carba NP (MCNP) test were used to screen for carbapenemase activity (15, 16). All strains were screened using real-time PCR for the presence of the following β -lactamase-encoding genes: *bla*_{KPC}, *bla*_{NDM}, *bla*_{VIM}, *bla*_{IMP}, *bla*_{OXA-48}, *bla*_{CTX-M}, *bla*_{TEM}, and *bla*_{SHV} (17–19). Strains positive by real-time PCR were subjected to standard PCR and sequencing (18, 20). Multilocus sequence typing (MLST) was carried out for carbapenemase-producing strains with the PubMLST scheme (<http://pubmlst.org/ecloacae/>).

Twelve *Enterobacteriaceae* isolates were obtained from 12 representative colonies, and strains of the following species were identified: *Klebsiella oxytoca*, *Citrobacter freundii*, *Enterobacter cloacae*, and *Citrobacter amalonaticus* (Table 1). The isolation from hospital cockroaches of such *Enterobacteriaceae* species with different levels of antimicrobial resistance has already been described (21). Additionally, in this study, we report for the first time the identification of the *Enterobacteriaceae* species *Citrobacter farmeri*, *Citrobacter koseri*, and *Enterobacter kobei* in German cockroaches. The majority of strains were isolated from the internal organs (Table 1), which could be explained by the great adaptation of these organisms in the intestinal environment (22). The antimicrobial susceptibility test results are presented in Table 1. The DDST was positive for 11 of 12 isolates. The MHT and the MCNP test were positive for only one *E. cloacae* isolate. Eleven of the 12 obtained strains harbored *bla*_{TEM}, and 75% ($n = 9$) of all isolates harbored the *bla*_{CTX-M-15} ESBL gene. One *E. cloacae* isolate belonging to sequence type 528 (ST528) harbored the *bla*_{OXA-48}, *bla*_{CTX-M-15}, and *bla*_{TEM} genes. According to the antibiogram results, some of the strains could express other β -lactamases not sought in this study. To our knowledge, this is the first description of CTX-M-15-producing *Enterobacteriaceae* isolated from *B. germanica* cockroaches. Few reports describing the presence of ESBL-producing *Enterobacteriaceae* in hospital cockroaches have been published (23–26). The most interesting finding in the present study was the first detection of OXA-48-producing *E. cloacae* in *B. germanica*. Since its first description in Turkey (27), numerous reports have indicated the spread of OXA-48 type carbapenemase in Mediterranean countries (28–30), where it represents the most common carbapenemase type (31). Subsequently, several studies have reported the isolation of OXA-48-producing *Enterobacteriaceae* from different samples (32, 33). Prior to its first description in Algeria, Poirel et al. suggested that OXA-48 may be endemic in this country (34). The huge spread of CTX-M-15 ESBL-producing *Enterobacteriaceae* and the emergence of OXA-48-type carbapenemase-producing *Enterobacteriaceae* in Algerian hospitals may explain the carriage of such drug-resistant bacteria by hospital cockroaches through their particular feeding habits (3, 4, 35).

The important role of cockroaches as a potential reservoir of multidrug-resistant bacteria demonstrated in this study and in earlier investigations with their presence in hospital environments, which has already been reported in Algeria (36) and other countries such as Turkey (9), Japan (37), and Cuba (38), urgently call for regulations regarding their control and elimination from clinical areas (11).

TABLE 1 Results of species identification and antibacterial susceptibility tests and β -lactamase determinants

Strain code	Source/ location ^a	Species identification	Antibiotic resistance ^b												β -lactamase-encoding genes			
			AMX	AMC	FOX	CTX	CAZ	FEP	ATM	ETP	IPM	TOB	GEN	AMK	CIP	SXT	TGC	CST
S1	IO/K	<i>Citrobacter amalonaticus</i>	R	S	R	R	S	S	S	I	R	S	S	R	S	S	S	<i>bla</i> _{CTX-M-15} , <i>bla</i> _{TEM}
S2	ES/P	<i>Citrobacter amalonaticus</i>	R	R	R	S	R	S	S	R	R	S	S	R	S	S	S	<i>bla</i> _{TEM}
S3	IO/K	<i>Citrobacter farmeri</i>	R	R	R	S	R	S	R	R	R	S	S	R	S	S	S	<i>bla</i> _{TEM}
S4	IO/T	<i>Citrobacter freundii</i>	R	R	R	R	R	R	R	R	R	S	R	S	S	S	S	<i>bla</i> _{CTX-M-15} , <i>bla</i> _{TEM}
S5	IO/K	<i>Citrobacter koseri</i>	R	S	S	R	R	R	R	S	I	R	S	S	R	S	S	<i>bla</i> _{CTX-M-15} , <i>bla</i> _{TEM}
S6	IO/T	<i>Enterobacter cloacae</i>	R	R	R	R	R	R	R	R	R	S	I	S	S	S	S	<i>bla</i> _{OXA-48} , <i>bla</i> _{CTX-M-15} , <i>bla</i> _{TEM}
S7	IO/T	<i>Enterobacter cloacae</i>	R	R	R	R	R	R	R	R	R	R	S	I	R	S	S	<i>bla</i> _{CTX-M-15} , <i>bla</i> _{TEM}
S8	ES/P	<i>Enterobacter cloacae</i>	R	R	R	R	R	R	R	S	R	S	S	S	S	S	S	<i>bla</i> _{CTX-M-15} , <i>bla</i> _{TEM}
S9	EST	<i>Enterobacter kobei</i>	R	R	R	R	R	S	R	R	S	S	S	S	S	S	S	None
S10	IO/T	<i>Klebsiella oxytoca</i>	R	S	R	R	R	R	R	S	S	R	S	S	R	S	S	<i>bla</i> _{CTX-M-15} , <i>bla</i> _{TEM}
S11	IO/K	<i>Klebsiella oxytoca</i>	R	S	S	R	R	I	I	S	S	R	S	S	S	S	S	<i>bla</i> _{CTX-M-15} , <i>bla</i> _{TEM}
S12	IO/K	<i>Klebsiella oxytoca</i>	R	S	S	R	R	R	S	S	R	R	S	S	S	S	S	<i>bla</i> _{CTX-M-15} , <i>bla</i> _{TEM}

^a ES, external surface; IO, internal organs; K, kitchen; P, patient's room; T, treatment room.

^b AMX, amoxicillin; AMC, amoxicillin-clavulanic acid; FOX, cefotaxime; CAZ, ceftazidime; FEP, cefepime; ATM, aztreonam; ETP, ertapenem; IPM, imipenem; TOB, tobramycin; GEN, gentamicin; CIP, ciprofloxacin; SXT, trimethoprim-sulfamethoxazole; TGC, tigecycline; CST, colistin; R, resistant; S, susceptible; I, intermediate.

ACKNOWLEDGMENTS

We are very grateful to Linda Hadjadj for technical assistance and to Mohamed Islam Hatcha for his contribution.

This work was partly funded by the Centre National de la Recherche Scientifique (CNRS 7278) and IHU Méditerranée Infection.

We declare no conflicts of interest.

REFERENCES

- Doi Y, Paterson DL. 2015. Carbapenemase-producing *Enterobacteriaceae*. *Semin Care Med* 36:74–84. <http://dx.doi.org/10.1055/s-0035-1544208>.
- Moxon CA, Paulus S. 2016. Beta-lactamases in *Enterobacteriaceae* infections in children. *J Infect* 72(Suppl):S41–S49. <http://dx.doi.org/10.1016/j.jinf.2016.04.021>.
- Cuzon G, Bentchouala C, Vogel A, Héry M, Lezzar A, Smati F, Doret L, Naas T. 2015. First outbreak of OXA-48-positive carbapenem-resistant *Klebsiella pneumoniae* isolates in Constantine, Algeria. *Int J Antimicrob Agents* 46:725. <http://dx.doi.org/10.1016/j.ijantimicag.2015.08.005>.
- Baba Ahmed-Kazi Tani Z, Arlet G. 2014. News of antibiotic resistance among Gram-negative bacilli in Algeria. *Pathol Biol* 62:169–178. <http://dx.doi.org/10.1016/j.patbio.2014.01.005>.
- Palmore TN, Henderson DK. 2013. Managing transmission of carbapenem-resistant *Enterobacteriaceae* in healthcare settings: a view from the trenches. *Clin Infect Dis* 57:1593–1599. <http://dx.doi.org/10.1093/cid/cit531>.
- Alrabaa SF, Nguyen P, Sanderson R, Baluch A, Sandin RL, Kelker D, Karlapalem C, Thompson P, Sams K, Martin S, Montero J, Greene JN. 2013. Early identification and control of carbapenemase-producing *Klebsiella pneumoniae*, originating from contaminated endoscopic equipment. *Am J Infect Control* 41:562–564. <http://dx.doi.org/10.1016/j.ajic.2012.07.008>.
- Akinjogunla OJ, Odeyemi AT, Udoinyang EP. 2012. Cockroaches (*Periplaneta americana* and *Blattella germanica*): reservoirs of multi drug resistant (MDR) bacteria in Uyo, Akwa Ibom State. *Sci J Biol Sci* 1:19–30.
- Kassiri H, Kassiri A, Kazemi S. 2014. Investigation on American cockroaches medically important bacteria in Khorramshahr hospital, Iran. *Asian Pacific J Trop Dis* 4:201–203. [http://dx.doi.org/10.1016/S2222-1808\(14\)60505-3](http://dx.doi.org/10.1016/S2222-1808(14)60505-3).
- Kutrup B. 2003. Cockroach infestation in some hospitals in Trabzon, Turkey. *Turk J Zool* 27:73–77.
- Brown C, Alhassan AN. 2015. Multiple-antibiotic-resistant bacteria from cockroaches trapped from a public hospital and a nearby students' hostel in Accra, Ghana. *Int J Biol Chem Sci* 8:1859–1864. <http://dx.doi.org/10.4314/ijbcs.v8i4.44>.
- Tilahun B, Worku B, Tachbele E, Terefe S, Kloos H, Legesse W. 2012. High load of multi-drug resistant nosocomial neonatal pathogens carried by cockroaches in a neonatal intensive care unit at Tikur Anbessa specialized hospital, Addis Ababa, Ethiopia. *Antimicrob Resist Infect Control* 1:12. <http://dx.doi.org/10.1186/2047-2994-1-12>.
- Vahabi A, Rafinejad J, Mohammadi P, Biglarian F. 2007. Regional evaluation of bacterial contamination in hospital environment cockroaches. *Iranian J Environ Health Sci Eng* 4:57–60.
- Seng P, Drancourt M, Gouriet F, La Scola B, Fournier P-E, Rolain JM, Raoult D. 2009. Ongoing revolution in bacteriology: routine identification of bacteria by matrix-assisted laser desorption ionization time-of-flight mass spectrometry. *Clin Infect Dis* 49:543–551. <http://dx.doi.org/10.1086/600885>.
- Jarlier V, Nicolas MH, Fournier G, Philippon A. 1988. Extended broad-spectrum β-lactamases conferring transferable resistance to newer β-lactam agents in *Enterobacteriaceae*: hospital prevalence and susceptibility patterns. *Rev Infect Dis* 10:867–878. <http://dx.doi.org/10.1093/clinids/10.4.867>.
- Lee K, Ki C, Yong D, Hoon S, Hwa J, Hee Y, Docquier J, Chong Y. 2010. Improved performance of the modified Hodge test with MacConkey agar for screening carbapenemase-producing Gram-negative bacilli. *J Microbiol Methods* 83:149–152. <http://dx.doi.org/10.1016/j.mimet.2010.08.010>.
- Bakour S, Garcia V, Loucif L, Brunel JM, Gharout-Sait A, Touati A, Rolain JM. 2015. Rapid identification of carbapenemase-producing *Enterobacteriaceae*, *Pseudomonas aeruginosa*, and *Acinetobacter baumannii* using a modified Carba NP test. *New Microbes New Infect* 7:89–93. <http://dx.doi.org/10.1016/j.nmni.2015.07.001>.
- Roschanski N, Fischer J, Guerra B, Roesler U. 2014. Development of a multiplex real-time PCR for the rapid detection of the predominant beta-lactamase genes CTX-M, SHV, TEM and CIT-type AmpCs in *Enterobacteriaceae*. *PLoS One* 9:e100956. <http://dx.doi.org/10.1371/journal.pone.0100956>.
- Mesli E, Berrazeg M, Drissi M, Bekkhoucha SN, Rolain JM. 2013. Prevalence of carbapenemase-encoding genes including New Delhi metallo-β-lactamase in *Acinetobacter* species, Algeria. *Int J Infect Dis* 17:e739–e743. <http://dx.doi.org/10.1016/j.ijid.2013.02.024>.
- Mellouk FZ, Bakour S, Meradji S, Al-Bayssari C, Bentakouk MC, Zouyed F, Djahoudi A, Boutefnouchet N, Rolain JM. 17 June 2016. First detection of VIM-4-producing *Pseudomonas aeruginosa* and OXA-48-producing *Klebsiella pneumoniae* in Northeastern (Annaba, Skikda) Algeria. *Microb Drug Resist* (Epub ahead of print.)
- Aubert D, Naas T, Heritier C, Poirel L, Nordmann P. 2006. Functional characterization of IS1999, an IS4 family element involved in mobilization and expression of beta-lactam resistance genes. *J Bacteriol* 188:6506–6514. <http://dx.doi.org/10.1128/JB.00375-06>.
- Elgderi RM, Ghengesh KS, Berbush N. 2006. Carriage by the German cockroach (*Blattella germanica*) of multiple-antibiotic-resistant bacteria that are potentially pathogenic to humans, in hospitals and households in Tripoli, Libya. *Ann Trop Med Parasitol* 100:55–62. <http://dx.doi.org/10.1179/13645906X78463>.
- Xue FU, Lefu YE, Feng GE. 2009. Habitat influences on diversity of bacteria found on German cockroach in Beijing. *J Environ Sci (China)* 21:249–254. [http://dx.doi.org/10.1016/S1001-0742\(08\)62259-7](http://dx.doi.org/10.1016/S1001-0742(08)62259-7).
- Cotton MF, Wasserman E, Pieper CH, Theron DC, Van Tubbergh D, Campbell G, Fang FC, Barnes J. 2000. Invasive disease due to extended spectrum beta-lactamase-producing *Klebsiella pneumoniae* in a neonatal unit: the possible role of cockroaches. *J Hosp Infect* 44:13–17. <http://dx.doi.org/10.1053/jhin.1999.0650>.
- Czajka E, Pancer K, Kochman M, Gliniewicz A, Sawicka B, Rabczenko D, Stypulkowska-Misiurewicz H. 2003. Characteristics of bacteria isolated from body surface of German cockroaches caught in hospitals. *Przegl Epidemiol* 57:655–662.
- Al-Marjani MF. 2008. Study of extended-spectrum β-lactamases producing enterobacteria isolated from cockroaches (*Periplaneta americana*) of Central Medicine City Hospital. *Al-Mustansiriya J Sci* 19:9–16.
- Doosti A, Pourabbas M, Arshi A, Chehelgerdi M, Kabiri H. 2015. TEM and SHV genes in *Klebsiella pneumoniae* isolated from cockroaches and their antimicrobial resistance pattern. *Osong Public Health Res Perspect* 6:3–8. <http://dx.doi.org/10.1016/j.phrp.2014.10.011>.
- Poirel L, Héritier C, Toltün V, Nordmann P. 2004. Emergence of oxacillinase-mediated resistance to imipenem in *Klebsiella pneumoniae*. *Antimicrob Agents Chemother* 48:15–22. <http://dx.doi.org/10.1128/AAC.48.1.15-22.2004>.
- Cuzon G, Ouanich J, Gondret R, Naas T, Nordmann P. 2011. Outbreak of OXA-48-positive carbapenem-resistant *Klebsiella pneumoniae* isolates in France. *Antimicrob Agents Chemother* 55:2420–2423. <http://dx.doi.org/10.1128/AAC.01452-10>.
- Potron A, Poirel L, Bussy F, Nordmann P. 2011. Occurrence of the carbapenem-hydrolyzing β-lactamase gene blaOXA-48 in the environment in Morocco. *Antimicrob Agents Chemother* 55:5413–5414. <http://dx.doi.org/10.1128/AAC.05120-11>.
- Kraiem AG, Zorgani A, Elahmer O, Hammami A, Chaaben BM, Ghengesh KS. 2015. New Delhi metallo-beta-lactamase and OXA-48 carbapenemases in Gram-negative bacilli isolates in Libya. *Libyan J Med* 10:29206.
- Djahmi N, Dunyach-Remy C, Pantel A, Dekhil M, Sotto A, Lavigne JP. 2014. Epidemiology of carbapenemase-producing *Enterobacteriaceae* and *Acinetobacter baumannii* in Mediterranean countries. *Biomed Res Int* 2014:1–11. <http://dx.doi.org/10.1155/2014/305784>.
- Al Bayssari C, Olaitan AO, Dabbousi F, Hamze M, Rolain JM. 2015. Emergence of OXA-48-producing *Escherichia coli* clone ST38 in fowl. *Antimicrob Agents Chemother* 59:745–746. <http://dx.doi.org/10.1128/AAC.03552-14>.
- Galler H, Feierl G, Petternel C, Reinthaler FF, Haas D, Grisold AJ, Luxner J, Zarfel G. 2014. KPC-2 and OXA-48 carbapenemase-harboring *Enterobacteriaceae* detected in an Austrian wastewater treatment plant. *Clin Microbiol Infect* 20:O132–O134. <http://dx.doi.org/10.1111/1469-0691.12336>.
- Poirel L, Potron A, Nordmann P. 2012. OXA-48-like carbapenemases: the phantom menace. *J Antimicrob Chemother* 67:1597–1606. <http://dx.doi.org/10.1093/jac/dks121>.
- Fakoorziba MR, Shahriari-Namadi M, Moemenbellah-Fard MD, Ha-

- tam GR, Azizi K, Amin M, Motavasel M. 2014. Antibiotics susceptibility patterns of bacteria isolated from American and German cockroaches as potential vectors of microbial pathogens in hospitals. *Asian Pacific J Trop Dis* 4:S790–S794. [http://dx.doi.org/10.1016/S2222-1808\(14\)60728-3](http://dx.doi.org/10.1016/S2222-1808(14)60728-3).
36. Menasria T, Moussa F, El-Hamza S, Tine S, Megri R, Chenchouni H. 2014. Bacterial load of German cockroach (*Blattella germanica*) found in hospital environment. *Pathog Glob Health* 108:141–147. <http://dx.doi.org/10.1179/204773214Y.0000000136>.
37. Saitou K, Furuhata K, Kawakami Y, Fukuyama M. 2009. Isolation of *Pseudomonas aeruginosa* from cockroaches captured in hospitals in Japan, and their antibiotic susceptibility. *Biocontrol Sci* 14:155–159. <http://dx.doi.org/10.4265/bio.14.155>.
38. Oliva GR, Diaz C, Fuentes GO, Martinez MD, Fernandez C, Cordovi R, Lago PM, Herrera N. 2010. *Blatella germanica* as a possible cockroach vector of micro-organisms in a hospital. *J Hosp Infect* 74:93–95. <http://dx.doi.org/10.1016/j.jhin.2009.09.002>.